

From band insulator to Mott insulator in one dimension

Michele Fabrizio^{1,3}, Alexander O. Gogolin² and Alexander A. Nersesyan^{3,4}

¹*International School for Advanced Studies and INFM, Via Beirut 4, 34014 Trieste, Italy*

²*Department of Mathematics, Imperial College, 180 Queen's Gate, London SW7 2BZ, United Kingdom*

³*The Abdus Salam International Centre for Theoretical Physics, P.O.Box 586, 34100, Trieste, Italy*

⁴*Institute of Physics, Tamarashvili 6, 380077, Tbilisi, Georgia*

(Draft: February 1, 2008)

We derive the phase diagram for the one-dimensional model of a ferroelectric perovskite recently introduced by Egami, Ishihara and Tachiki [*Science*, **261**, 1307 (1993)]. We show that the interplay between covalency, ionicity and strong correlations results in a spontaneously dimerized phase which separates the weak-coupling band insulator from the strong-coupling Mott insulator. The transition from the band insulator to the dimerized phase is identified as an Ising critical point. The charge gap vanishes at this single point with the optical conductivity diverging as $\sigma(\omega) \sim \omega^{-3/4}$. The spin excitations are gapless above the second transition to the Mott insulator phase.

71.30.+h, 77.80.-e, 71.10.Pm

Much effort have been devoted to understanding the response of strongly correlated electron systems to lattice distortions. Generally, one expects that the tendency to a charge density wave instability modulated by a lattice distortion is increased by (repulsive) interactions. A surprisingly pronounced enhancement has recently been observed in numerical studies of a prototype one dimensional (1D) model for ferroelectric perovskites, proposed by Egami, Ishihara and Tachiki [1].

The Hamiltonian of their model is

$$\hat{H} = \sum_{i\sigma} \left[-t \left(c_{i\sigma}^\dagger c_{i+1\sigma} + \text{H.c.} \right) + \Delta (-1)^i n_{i\sigma} \right] + U \sum_i n_{i\uparrow} n_{i\downarrow}, \quad (1)$$

at half-filling. The odd and even sites represent oxygen atoms (O) and a generic cation (C), respectively, with the energy difference $E_C - E_O = 2\Delta$. The hopping amplitude t determines the amount of covalency and U stands for the on-site Coulomb repulsion. In spite of its apparent simplicity, model (1) reveals nontrivial physics. At $U = 0$ it describes a band insulator (BI) of a mixed ionic-covalent character, with a spectral gap for all excitations. At $\Delta = 0$ it is a Mott insulator (MI) with a finite charge gap, m_c , but with gapless spin excitations. In fact, in the strong-coupling limit $U \gg (t, \Delta)$, (1) can be mapped onto a Heisenberg model by projecting out all doubly occupied sites. At the leading order, only a nearest neighbour exchange $J = 4t^2U/(U^2 - 4\Delta^2)$ is generated. The resulting Heisenberg chain is well known to possess a gapless spin spectrum. The issue of interest is

the nature of the crossover from the BI regime to the MI regime which, on general grounds, is expected to occur in the region where the single-particle BI gap Δ becomes comparable with the MI charge gap m_c .

Thus, increasing U at fixed Δ , one expects a transition to a spin-gapless phase. In addition to this *spin transition*, the charge degrees of freedom should be also affected in the course of the crossover. This can easily be understood in the case $t \ll (U, \Delta)$ where, upon increasing U , the system approaches a mixed-valence regime in which the two charge configurations, $\text{O}^{-2}\text{C}^{+2}$ with energy $U - 2\Delta$ and $\text{O}^{-1}\text{C}^{+1}$ with zero energy, become degenerate. Therefore, those excitations responsible for the charge redistribution among these two configurations should soften around some value of U . The finite-size simulations [1–5] suggest that the mixed-valence regime is accompanied by a strongly enhanced response to the coupling with zone-centre optical phonons ξ ,

$$\hat{H}_{ep} = -\lambda \xi \sum_i \mathcal{D}_i, \quad \mathcal{D}_i = \sum_{\sigma} (-1)^i \left(c_{i\sigma}^\dagger c_{i+1\sigma} + \text{H.c.} \right), \quad (2)$$

where λ is the electron-phonon coupling constant and $\mathcal{D}_i$ is the dimerization (or staggered bond-density) operator. In fact, the so-called average dynamical charge $Z_* = \lim_{\xi \rightarrow 0} (P(\xi) - P(0))/\xi$, where $P(\xi)$ is the polarization, diverges at a particular point, even in finite-size simulations [4,3]. This is due to the *accidental* degeneracy between two singlet ground states with opposite parity which occurs in any finite-size system with the number of sites multiple of 4 [4,3]. More importantly, Resta and Sorella [5] have recently observed that the localization length of the ground-state wave function diverges at some value of U , which suggests vanishing of the charge gap; hence the *charge transition*.

However, the overall picture which emerges from numerical simulations and the existing mean-field calculations [3], clearly underestimating the crucial role of fluctuations in 1D, is still far from being satisfactory. It remains unclear whether the charge instability occurs at the same point as the spin transition. Even the nature of the transition (transitions?) is not known. Is it a first-order transition or is it a second-order one? In the latter case, what is the universality class it belongs to? To the best of our knowledge, there have been no consistent analytic attempts to resolve these intriguing questions.

In this Letter we present theoretical arguments showing that there are two continuous transitions: the spin transition of the Kosterlitz-Thouless (KT) type at $U = U_{c2}$, and the charge transition at $U = U_{c1} < U_{c2}$, the latter identified as an Ising critical point where the charge gap vanishes. In the intermediate region, $U_{c1} < U < U_{c2}$, the site-parity is spontaneously broken, and the system is characterised by a doubly degenerate, dimerized ground state, with $\langle \mathcal{D}_i \rangle$ being the order parameter.

We start our discussion with addressing the mechanism of the spin gap generation in the MI phase. For the single-chain Hamiltonian (1), which is $SU(2)$ and site-parity invariant, the only possibility compatible with the symmetries is a spontaneous spin-dimerization, like the one occurring at $J_2/J_1 \simeq 0.24$ in the frustrated Heisenberg chain with nearest-neighbour (J_1) and next-nearest-neighbour (J_2) exchanges. The same scenario is realized in our case below a critical value U_{c2} which depends on Δ and t . This result agrees with the exact diagonalization of a 4-site chain where the next-to-nearest neighbour exchange starts to play a role. At large U , the ground state is an odd-parity singlet, and the lowest excitation is a spin-triplet, followed at higher energy by an even-parity singlet. As U decreases, first the even-parity singlet and the triplet cross, and finally the two opposite-parity singlets cross. For $(U, \Delta) \gg t$ these two level crossings occur at $U \simeq 2\Delta + 1.86t$ and at $U \simeq 2\Delta + t$, respectively (see also Ref. [3]). The same is observed in the J_1 - J_2 Heisenberg model when J_2 is increased [6]. The crossing of the triplet and singlet excited states in finite-size systems becomes the spontaneous spin-dimerization transition in the thermodynamic limit. The crossing of the two opposite-parity singlets, taking place at $J_2 = 0.5 J_1$, corresponds to the Majumdar-Gosh solvable limit [7].

The appearance of a spontaneously dimerized insulating (SDI) phase can be also inferred starting from the BI phase. In this limit we need an improved version of the mean-field calculation presented in Ref. [3]. We introduce a two component Fermi field $\Psi_{k\sigma}^\dagger = (c_{k\sigma}^\dagger, c_{k+\pi\sigma}^\dagger)$, $k \in [-\pi/2, \pi, 2]$, and the operators

$$\hat{\tau}_{a\sigma}(q) = \sum_k \Psi_{k\sigma}^\dagger \tau_a \Psi_{k+q\sigma}, \quad (a = 0, 1, 2, 3), \quad (3)$$

where τ_a are the Pauli matrices including the unit matrix τ_0 . The staggered charge/spin densities correspond to $\hat{\tau}_{1\uparrow}(0) \pm \hat{\tau}_{1\downarrow}(0)$, while the dimerization is $\hat{\tau}_{2\uparrow}(0) + \hat{\tau}_{2\downarrow}(0) \equiv \sum_i \mathcal{D}_i$ [see Eq.(2)]. A generic interaction compatible with the symmetry of (1) can be written as

$$\begin{aligned} \hat{H}_{int} = & \frac{1}{4L} \sum_{q, \sigma, \sigma'} [(g_1 + g_3) \hat{\tau}_{1\sigma}(q) \hat{\tau}_{1\sigma'}(-q) \\ & + (g_1 - g_3) \hat{\tau}_{2\sigma}(q) \hat{\tau}_{2\sigma'}(-q) + 2g_2 \hat{\tau}_{0\sigma}(q) \hat{\tau}_{0\sigma'}(-q)]. \end{aligned} \quad (4)$$

The coupling constants are written using the g -ology jargon [8]: g_1, g_2 and g_3 label the backward, forward and

Umklapp scattering amplitudes, respectively. In the special case of (1), $g_1 = g_2 = g_3 = U$, and the $\hat{\tau}_2$ - $\hat{\tau}_2$ term vanishes. Therefore, in any mean-field treatment of the original Hamiltonian, only variational states with finite average values of $\hat{\tau}_{1\sigma}(0)$, i.e. the staggered charge and spin densities, can be explored. However, the equality among the g 's is already lost at the first order of perturbation theory. More refined weak-coupling schemes, such as renormalization group [8] or the bosonization technique [10], show that renormalization occurring at energies well below the bandwidth cutoff but still larger than $\max(\Delta, m_c)$ enhances g_3 and g_2 but suppresses g_1 . This implies generation of an effective coupling $(g_3 - g_1)$ which may lead to the appearance of a finite average value of $\hat{\tau}_2(0)$, in addition to the average $\langle \hat{\tau}_1(0) \rangle$ induced by the Δ -term. By including these renormalization effects within the leading logarithmic approximation [9], we indeed find a sequence of two transitions. First, at $U = U_{c1}$, a finite average $\langle \hat{\tau}_2(0) \rangle$ continuously appears. At a larger $U = U_{c2}$ a triplet exciton gets soft signalling the transition to the MI phase. Although these two transitions are physically distinct within this scheme, to estimate the difference between U_{c1} and U_{c2} one has to go beyond the leading logarithmic approximation, which is not feasible. For this reason, we can only safely state that $U_{c1} < U_{c2}$, with both critical points being of the order $2\pi t / \ln(t/\Delta)$.

In what follows, we develop the *low-energy* effective field theory for the lattice model (1). Considering the weak-coupling case, $(U, \Delta) \ll t$, we linearise the spectrum and pass to the continuum limit by substituting $a_0^{-1/2} c_{n\sigma} \rightarrow i^n \psi_{R\sigma}(x) + (-i)^n \psi_{L\sigma}(x)$, $x = na_0$, where a_0 is the lattice spacing, and $\psi_{R,L}(x)$ are the right and left components of the Fermi field. These fields can be bosonized in a standard way [10]: $\psi_{R,L;\sigma} = (2\pi a_0)^{-1/2} e^{\pm i\sqrt{4\pi}\phi_{R,L;\sigma}}$, where $\phi_{R(L),\sigma}$ are the right(left)-moving Bose fields. We define $\Phi_\sigma = \phi_{R\sigma} + \phi_{L\sigma}$ and introduce linear combinations, $\Phi_c = (\Phi_\uparrow + \Phi_\downarrow)\sqrt{2}$ and $\Phi_s = (\Phi_\uparrow - \Phi_\downarrow)\sqrt{2}$, to describe the charge and spin degrees of freedom, respectively. Then the Hamiltonian density of the bosonized model is given by: $\mathcal{H}_{eff} = \mathcal{H}_c + \mathcal{H}_s + \mathcal{H}_{cs}$. Here the charge and spin sectors are described by

$$\begin{aligned} \mathcal{H}_c = & \frac{v_c}{2} [\Pi_c^2 + (\partial_x \Phi_c)^2] \\ & - \frac{g_3}{\pi^2 a_0} \cos \sqrt{8\pi} \Phi_c + \frac{2g_2 a_0}{\pi} \partial_x \phi_{cR} \partial_x \phi_{cL} \\ \mathcal{H}_s = & \frac{v_s}{2} [\Pi_s^2 + (\partial_x \Phi_s)^2] \\ & + \frac{g_1}{\pi^2 a_0} \cos \sqrt{8\pi} \Phi_s - \frac{2g_1 a_0}{\pi} \partial_x \phi_{sR} \partial_x \phi_{sL} \end{aligned} \quad (5) \quad (6)$$

where $\Pi_{c,s}$ are the momenta conjugate to $\Phi_{c,s}$, and $v_{c,s}$ are the velocities of the charge and spin excitations. Actually, $\mathcal{H}_s$ in (6) is an Abelian bosonized version of the $SU(2)_1$ -symmetric Wess-Zumino-Novikov-Witten model with a marginally irrelevant ($g_1 > 0$) current-current per-

turbation, $-g_1 \mathbf{J}_R \cdot \mathbf{J}_L$, given by the last two terms in (6). The two sectors of the theory are coupled by the Δ -term:

$$\mathcal{H}_{cs} = -(2\Delta/\pi a_0) \sin \sqrt{2\pi} \Phi_c \cos \sqrt{2\pi} \Phi_s. \quad (7)$$

In fact, the effective Hamiltonian $\mathcal{H}_{eff}$ represents the low-energy limit for a wide class of lattice models including (1). Such lattice effects as nearest-neighbour repulsion, or a difference between the Coulomb repulsion on the O-atoms and the C-atoms, will only renormalize the parameters of $\mathcal{H}_{eff}$, in particular the charge stiffness constant (K_c) and the velocities $v_{c,s}$.

Starting from the MI phase, one can firmly justify the opening of a spin gap and spontaneous dimerization upon decreasing U . Assuming $U \gg \Delta$, in which case the spin sector is gapless while the charge one is gapped, one can integrate out the charge degrees of freedom to obtain at the second order in Δ a nontrivial renormalization of g_1 : $\tilde{g}_1 = g_1 - (\Delta/m_c)^2 (v_c/a_0)$. Until $\tilde{g}_1 > 0$, the spin spectrum remains gapless. However, when U is decreased, $\tilde{g}_1$ eventually becomes negative, and the system undergoes a continuous (KT) transition to the dimerized phase with a nonzero spin gap. Using the exact result for the small- U Hubbard model, $m_c \sim \sqrt{U} t e^{-2\pi t/U}$, one concludes that

$$U_{c2} = \frac{2\pi t}{\ln(t/\Delta)} \left[1 + O\left(\frac{\ln \ln(t/\Delta)}{\ln(t/\Delta)}\right) \right], \quad (8)$$

in agreement with the previous estimate.

Bosonization also offers a simple understanding of the region $U < U_{c2}$. In what follows, we shall focus on the nature of the transition at $U = U_{c1}$. To this end, the Hamiltonian $\mathcal{H}_{eff}$, given by (5)-(7), will be regarded as a *phenomenological* Landau-Ginzburg energy functional, in the sense that all the couplings are effective ones obtained by integrating out high-energy degrees of freedom.

We start with looking at the saddle points of the effective potential in $\mathcal{H}_{eff}$ which obey the equations (here $\varphi_{c,s} = \sqrt{2\pi} \Phi_{c,s}$)

$$\cos \varphi_c \left[4\tilde{g}_3 \sin \varphi_c - 2\pi \tilde{\Delta} \cos \varphi_s \right] = 0 \quad (9)$$

$$\sin \varphi_s \left[2\pi \tilde{\Delta} \sin \varphi_c - 4\tilde{g}_1 \cos \varphi_s \right] = 0, \quad (10)$$

supplemented by stability conditions. At $\tilde{g}_3 < \pi \tilde{\Delta}/2$ one finds two sets of minima (defined modulo 2π) located at $\varphi_s = 0$, $\varphi_c = \pi/2$, and $\varphi_s = \pi$, $\varphi_c = -\pi/2$. These sets characterise the BI phase. Indeed, the vacuum-vacuum transitions, $\Delta \varphi_{s(c)} = \pm\pi$, describe stable topological excitations carrying the charge $Q = \Delta \varphi_c / \pi = \pm 1$ and spin $S^z = \Delta \varphi_s / 2\pi = \pm 1/2$ and therefore coinciding with “massive” single-fermion excitations of the BI. The situation changes at $\tilde{g}_3 > \pi \tilde{\Delta}/2$ where each minimum in the charge sector splits into two degenerate minima and thus transforms to a local *double-well* potential. The new minima are given by $\varphi_s = 0$, $\varphi_c = \phi_0$, $\pi - \phi_0$, and $\varphi_s = \pi$, $\varphi_c = -\phi_0$, $-\pi + \phi_0$, where $\phi_0 = \arcsin(\pi \tilde{\Delta} / 2 \tilde{g}_3)$. They

describe the SDI phase. In this region, the dimerization operator $\mathcal{D}(x) \sim \cos \varphi_c(x) \cos \varphi_s(x)$ acquires a finite expectation value. Notice that the location of the minima in the spin sector, and hence the spin quantum numbers of the topological excitations, are the same as in the BI phase ($\tilde{g}_3 < \pi \tilde{\Delta}/2$). Therefore the spin part of the spectrum formed in the SDI phase smoothly transforms into that of the BI phase. The transition at $\tilde{g}_3 = \pi \tilde{\Delta}/2$ mainly involves the charge degrees of freedom which undergo dramatic changes. In particular, the charge quantum numbers become *fractional*, depending on ϕ_0 . The Z_2 -degeneracy of the SDI state implies the existence of topological kinks carrying the spin $S = 1/2$ and charge $Q = \pm 2\varphi_0/2$. These massive excitations, described by interset vacuum-vacuum transition, interpolate between the neutral spinons of the MI phase and single-fermion excitations of the BI phase. However, the double-well local structure of the effective potential in the charge sector gives rise to *singlet* kink excitations as well, whose mass and charge $Q = 1 - 2\phi_0/\pi$ vanish at $\tilde{g}_3 = \pi \tilde{\Delta}/2$. On the BI side of the phase diagram, these singlet excitations appear as excitonic bound states inside the single-particle gap. Precisely at $\tilde{g}_3 = \pi \tilde{\Delta}/2$, the effective potential becomes $\sim \varphi_c^4$, which is well known to describe the Ising universality class.

We have checked that the SDI solution remains stable in the region

$$\sqrt{\tilde{g}_1 \tilde{g}_3} \leq \frac{\pi}{2} \tilde{\Delta} \leq \tilde{g}_3. \quad (11)$$

This condition can be satisfied if $\tilde{g}_1 < \tilde{g}_3$, which is indeed true for our phenomenological Hamiltonian. However, separation between the transition points U_{c1} and U_{c2} becomes even stronger when the electron-phonon interaction, always present in realistic situations, is taken into account. Let us assume that the zone-centre optical phonons are dispersionless, with a characteristic frequency ω_0 being the largest energy scale in the problem. In this limit, the phonons mediate an instantaneous interaction between the electrons which adds to the Hubbard repulsion and leads to a renormalization of the “bare” coupling constants: $g_3 \rightarrow g_3 + g_0$, $g_1 \rightarrow g_1 - g_0$, where $g_0 \sim \lambda^2/\omega_0$. The transition points U_{c1} and U_{c2} will start moving apart upon increasing g_0 , and if the electron-phonon coupling is large enough ($g_0 > g_1$), a finite spin gap will be generated even at $\Delta = 0$. Qualitatively the same situation occurs upon relaxing the condition of extremely quantum phonons (large ω_0) at a fixed λ . This observation explains that the two transitions can be, in principle, completely disentangled and, on the other hand, it provides an extra support to the fact that the charge transition separates the BI and SDI phases.

The Ising scenario at $U = U_{c1}$ can be more rigorously proven by studying the charge degrees of freedom with the spin bosonic field locked at $\Phi_s = 0$ or $\sqrt{\pi}/2$. Such approach will certainly be valid in a small vicinity of

U_{c1} because, as we already mentioned, the structure of the gapped spin sector is qualitatively the same both in the BI and SDI phases. The charge sector can then be described by a double sine-Gordon Hamiltonian

$$\hat{H}_c(x) = \frac{v_c}{2} \left[\Pi_c(x)^2 + (\partial \Phi_c(x))^2 \right] - \frac{g}{\pi^2 a_0} \cos \sqrt{8\pi K_c} \Phi_c(x) - \frac{2\Delta}{\pi a_0} \sin \sqrt{2\pi K_c} \Phi_c(x), \quad (12)$$

where, we stress once more, all the parameters are purely phenomenological, and part of the interaction has been absorbed in the rescaling parameter $K_c < 1$. A quantum phase transition of an Ising type in the double sine-Gordon model was recently discovered by Delfino and Mussardo [11]. In the whole parameter region of interest to us, i.e. $g > 0$ and $K_c < 1$, they argued that the existence of an Ising critical point is a universal property of this model. This allows us to investigate this model in the vicinity of the point $K_c = 1/2$, where the problem greatly simplifies. At this point, the Hamiltonian (12) can be mapped onto two coupled 2D Ising models, or equivalently, two coupled quantum Ising chains in a transverse magnetic field (a deformation of the Ashkin-Teller model). One of the two chains always remains off-critical, while the other one can pass through a critical point by fine tuning the parameters of the model. The details will be presented elsewhere [9]. Here we just quote our main findings, which can be easily understood once the Ising scenario is assumed. The most relevant perturbation at an Ising transition is the magnetic field h , which, in the ordered phase, selects one of the two equivalent vacua. By analogy, the role of h is played in our case by the explicit dimerization ξ , Eq.(2). The dimerization operator $\mathcal{D}$ (which also plays the role of the charge polarizaton operator) is therefore proportional to the order parameter σ of the quantum critical Ising model. The Ising mapping also allows us to identify the electron current operator, $j \sim \partial_t \sigma$, and the average electron charge operator $\rho \sim -\partial_x \sigma$. These findings are perfectly consistent with the general relations $\rho \sim -\partial_x \mathcal{D}$ and $\partial_t \rho + \partial_x j = 0$ (continuity) valid for insulators (i.e., in the absence of free carriers). From known results on the correlation functions at the Ising transition we conclude that the susceptibility to the explicit dimerization, namely the real part of the polarizability $\chi(\omega)$, diverges at low frequencies as $\omega^{-7/4}$, and the optical conductivity, $\sigma(\omega) \sim \omega^{-3/4}$ (no Drude peak), is ‘semi-metallic’.

In conclusion, we have identified the origin of the anomalous response to an electron-phonon coupling in the 1D ferroelectric perovskite model (1), which turns out to be far stronger than that of free electrons ($\Delta = U = 0$), of a band insulator ($U = 0$), or of a pure Hubbard model ($\Delta = 0$). Our findings are summarised in Fig.1, where the phase diagram is shown at fixed Δ and t as a function of U . Between the BI ($U < U_{c1}$) and MI ($U > U_{c2}$) phases there exists a spontaneously dimerized phase even

in the absence of an explicit electron-phonon interaction [12]. The transition from the MI to the spontaneously dimerized insulator is of the KT type, accompanied by opening of the spin gap. The transition from the dimerized phase to the BI has been shown to be of the Ising type, with the charge gap closing at this single point [12].

We acknowledge useful discussions with D. Edwards, Yu Lu, and E. Tosatti. We particularly thank G. Mussardo and S. Sorella for their enlightening comments. MF is partly supported by INFM, under project PRA HTSC. AOG is supported by the EPSRC of the United Kingdom.

-
- [1] T. Egami, S. Ishihara, and M. Tachiki, *Science* **261**, 1307 (1993).
 - [2] S. Ishihara, T. Egami, and M. Tachiki, *Phys. Rev. B* **49**, 8944 (1994); S. Ishihara, M. Tachiki, and T. Egami, *ibid.* **49**, 16123 (1994).
 - [3] G. Ortiz and R. Martin, *Phys. Rev. B* **49**, 14202 (1994); G. Ortiz, P. Ordejón, R. Martin, and G. Chiappe, *ibid.* **54**, 13515 (1996).
 - [4] R. Resta, and S. Sorella, *Phys. Rev. Lett.* **74**, 4738 (1995).
 - [5] R. Resta, and S. Sorella, *Phys. Rev. Lett.* **82**, 370 (1999).
 - [6] K. Okamoto and K. Nomura, *Phys. Lett. A* **169**, 433 (1993); I. Affleck, D. Gepner, H.J. Schulz, and T. Ziman, *J. Phys. A* **22**, 511 (1989).
 - [7] C.K. Majumdar and D.K. Ghosh, *J. Math. Phys.* **10**, 1388 (1969).
 - [8] I.E. Dzyaloshinskii and A.I. Larkin, *Sov. Phys. JETP* **34**, 422 (1972); J. Sólyom, *Adv. Phys.* **28**, 201 (1979).
 - [9] M. Fabrizio, A.O. Gogolin and A.A. Nersesyan, unpublished.
 - [10] See for instance A.O. Gogolin, A.A. Nersesyan, and A.M. Tsvelik, *Bosonisation and Strongly Correlated Systems*, Cambridge University Press, Cambridge (1998).
 - [11] G. Delfino and G. Mussardo, *Nucl. Phys. B* **516**, 674 (1998).
 - [12] Evidence in favour of a two transition scenario was recently uncovered numerically by N. Gidopoulos, S. Sorella and E. Tosatti (unpublished).




FIG. 1. Phase diagram of (1) as function of U at fixed Δ and t . $\Delta_{c(s)}$ stands for the charge (spin) gap, $\mathcal{D}_i$ is the dimerization operator, whose average is finite in the spontaneously dimerized insulator but vanishes both in the band and in the Mott insulators.